

EXPLANATION OF PLATE


Figs. 1a, 1b: *Crania antarcticaensis* HATAI, n. sp. $\times 2$. Inner (Fig. 1a) and outer (Fig. 1b) views of dorsal valve. Loc. Off Cape Cook Island, 700 m depth.

Figs. 2a, 2b: *Crania antarcticaensis* HATAI, n. sp. $\times 2$. Inner (Fig. 2a) and outer (Fig. 2b) views of dorsal valve. Loc. Off Cape Cook Island, 500 m depth.

Figs. 3a, 3b: *Crania antarcticaensis* HATAI, n. sp. $\times 2$. Inner (Fig. 3a) and outer (Fig. 3b) views of dorsal valve. Loc. Off Cape Cook Island, 500 m depth.

Figs. 4a, 4b: *Crania antarcticaensis* HATAI, n. sp. $\times 2$. Inner (Fig. 4a) and outer (Fig. 4b) views of dorsal valve. Loc. Gunnerus Bank, 590 m depth.

Figs. 5a, 5b: *Crania antarcticaensis* HATAI, n. sp. $\times 2$. Inner (Fig. 5a) and outer (Fig. 5b) views of dorsal valve. Loc. Gunnerus Bank, 590 m depth.



HATAI: *Crania* (Brachiopoda) from Antarctica.